



Investigations of beer carbohydrates with increased ^{13}C sensitivity

Sundekilde, Ulrik K.; Meier, Sebastian

Publication date:
2016

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Sundekilde, U. K., & Meier, S. (2016). *Investigations of beer carbohydrates with increased ^{13}C sensitivity*. Abstract from EUROMAR 2016, Aarhus, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Investigations of beer carbohydrates with increased ^{13}C sensitivity

Ulrik K. Sundekilde,¹ Sebastian Meier²

¹Department of Food Science, Aarhus University, Denmark. ²Department of Chemistry, Technical University of Denmark, Denmark.

Complex carbohydrates are an integral part of foods and the structural assignment of these complex oligo- or polysaccharides is a necessary step for understanding their biological functions and biosynthetic pathways. NMR is a powerful and often used technique in this structure determination and it often relies on natural ^{13}C abundance, which comes with severe limitations – low sensitivity and highly overlapping spectra. Insight into the multifaceted mixture of different carbohydrates at natural ^{13}C isotopic abundance was achieved through ^{13}C projections from two-dimensional ^1H - ^{13}C HSQC experiments using high-resolution in the second dimension, thereby providing a highly detailed mapping of a central compound class that is not easily amenable to other detection modalities. Carbohydrate structures were investigated by use of the anomeric reporter signals (Figure 1A). Multivariate data analysis techniques were used to examine the carbohydrate profile of ale, lager and Lambic beer types to identify differences (Figure 1B), which could stem from differences in the fermentation processes. Moreover, the carbohydrate profiles are correlated to the beer alcohol content, thereby providing information about biological and biotechnological processes in beer production.




Figure 1. A) ^{13}C spectra created from positive projection of F1 in ^1H - ^{13}C HSQC spectra of 42 beers. b) S-line plot from an OPLS-DA model on ^{13}C projected spectra of beers (ale and lager).